Торіс	Evaluation of the 24-hour Pseudalert* test versus Standard Methods <sup>1</sup>		
	9213F and 9213E for the detection of <i>Pseudomonas aeruginosa</i> in		
	both public and private spa water samples		
Title	Poster title: "Incidence of Pseudomonas aeruginosa in Private Spa		
	Water"		
Author(s)	Nancy Hall, Cathy Lord, John Kempf, Carrie Lueck, Cindy Rieflin,		
	and Karen Owens; State Hygienic Laboratory, University of Iowa		
	Research Park, Iowa City, IA 52242		
Date	April 2011, poster presented at Iowa meeting of State Public Health		
	Labs and Officials		

#### Highlights:

- Twelve public spas from the Scott County area were tested in parallel by both new and current methods; additionally, 65 private individuals submitted water samples from their private spas.
- Spa chemistry information (e.g. total chlorine/bromine, pH) was obtained at time of collection.
- For the public spa study there was only one positive *Pseudomonas aeruginosa* sample (1/12 or 8%) but for the private spa study there were significantly more positives (41/65 or 63%).
- The Pseudalert test was comparable to the *Standard Method* tests, demonstrating very low false positive and negative results with a sensitivity and specificity of 95.4% and 100%, respectively.
- 56% of samples with unacceptable chemistry levels and 4 % of samples with acceptable chemistry levels were positive for *Pseudomonas aeruginosa*. This has led the authors to both recommend public health intervention and develop strategies for spa waters.

1. *Standard Methods for the Examination of Water and Wastewater*, 21<sup>st</sup> Ed, 2005, APHA, AWWA & WEF, Washington, DC

\* Pseudalert is a trademarks or registered trademark of IDEXX Laboratories, Inc. or its affiliates in the United States and/or other countries





# Abstract

Background: Pseudomonas aeruginosa spa testing is a good indicator of overall spa maintenance. If this organism is present in the water, there is a potential health risk for the bather including skin rash or other complications. Public spa facilities are required by state law to collect a monthly water sample but privately-owned spas/whirlpools are NOT required to test their water for this organism.

**Purpose:** The original purpose of this study was to validate a faster Pseudomonas aeruginosa method (24-hr versus the current 48 to 96-hr test) that recently became available. The results not only demonstrated that the two methods were comparable but revealed a larger public health problem; that many private spa waters are positive for *Pseudomonas* aeruginosa indicating poor maintenance procedures and potential health hazard. This poster will hopefully start the dialog among the public health community as to how we can successfully address this problem and offer possible solutions and educational opportunities to the private sector and spa dealerships.

### Introduction

In 2010 the State Hygienic Laboratory (SHL) participated in a method validation study of a new 24-hr Pseudomonas aeruginosa test called Pseudalert<sup>™</sup> (IDEXX Laboratories, Inc., Westbrook, ME). The overall objective of this trial was to compare the performance of the new method to the current Standard Methods for the detection and quantification of *P*. aeruginosa in spa water samples. Twelve public spas from the Scott County area were tested in parallel by both new and current methods. Since more spa samples were needed for this study, a mass email was sent to University of Iowa faculty and staff asking for their participation. Sixty-five private individuals submitted water samples from their private spas for *Pseudomonas* testing. Some spa chemistry information (e.g. total chlorine/bromine, pH) was obtained at time of collection.

# Incidence of *Pseudomonas aeruginosa* in Private Spa Water

Nancy Hall, Cathy Lord, John Kempf, Carrie Lueck, Cindy Rieflin, and Karen Owens State Hygienic Laboratory, University of Iowa Research Park, Iowa City, IA 52242

# Methods

The three *Pseudomonas aeruginosa* methods performed in this study were the following: current SHL presence/absence test (Standard Method 9213F) performed on the public spa samples, membrane filter (MF) test (Standard Method 9213E) performed on the private spa samples, and the new Pseudalert<sup>™</sup> test performed on both sample sets.

## Results

Method Comparison: The combined results of the new Pseudalert<sup>™</sup> method and the two Standard Methods (both public and private spa sample set) are given in Table 1. The Pseudalert <sup>™</sup> method was comparable to the Standard Method tests by demonstrating very low false positive and negative results with a sensitivity and specificity of 95.4% and 100%, respectively.

#### **Pseudomonas and Spa Chemistry Results:**

For the public spa study there was only one positive Pseudomonas aeruginosa sample (1/12 or 8%) but for the private spa study there were significantly more positives; 41/65 or 63% as shown in Table 2. Twelve private spa samples also had mold present on the MF plate (with or without Pseudomonas) which also indicated unsanitary conditions. A high percentage of private positive Pseudomonas samples (56%) had "not acceptable" chemistry levels while only 4 (9%) samples had acceptable chemistries (using the preferred operating pH and chlorine/bromine levels stated in the lowa regulations (IAC 641-15) for public spas). When the spa chemistry was "not acceptable," it was primarily the chlorine/bromine levels that were out of range (Figure 1). It is not clear from the information provided whether some spas were using additional or other disinfection technologies such as silver-copper ionization, ozone, or UV at the time of sample collection.

### **Table 1. Pseudalert<sup>™</sup> and Both Standard** Methods Result Summary

	Pseudalert POSITIVE	Pseudalert NEGATIVE	Totals
Standard Methods POSITIVE	42	0	42
Standard Methods NEGATIVE	2*	33	35
Totals	44	33	77

\*Two samples SM 9213E only

Sensitivity =	42/44 = 9	5.4%
Specificity =	33/33 = 1	00%


PPV = 42/42 = 100%NPV = 33/35 = 94.3%

### Table 2. Private Spa Pseudomonas and **Chemistry Results**

<i>Pseudomonas</i> Results	Total	Acceptable Chemistries	Not Acceptable Chemistries*	No Chemistry information provided
Positive	<b>41 (63%)</b>	4	23 (56%)	14
Negative	24 (37%)	4	5	15

\*Not Acceptable Spa Chemistries: either chlorine/bromine <2 mg/L OR pH <7.2.

### Figure 1: Private Spa Chlorine/Bromine Levels for *Pseudomonas* Positive and **Negative Samples**



# Discussion

- > The public health community needs to address this public health problem and develop strategies to reduce the high incidence of *Pseudomonas* in private spa water.
- Possible public health strategies include the following:
- Recommend routine Pseudomonas aeruginosa monitoring for private spas
- Recommend testing should be done within the first month of operation and at least quarterly thereafter (or more frequently as necessary) to check effectiveness of spa sanitation practices
- Develop educational brochures and seminars for private individuals AND spa dealers regarding proper spa maintenance

# Conclusions

- ➤ The new Pseudalert<sup>™</sup> test by IDEXX Laboratories was comparable to current Standard Methods for Pseudomonas aeruginosa testing in both public and private spa samples.
- > This study demonstrated a high percentage (63%) of positive Pseudomonas aeruginosa private spa samples that may pose a health risk to bathers.
- Based on the high incidence of Pseudomonas aeruginosa in private spa water, a public health intervention is needed and strategies should be developed.

# Acknowledgements

- Dave Townsend and Clifford Young, IDEXX Laboratories, for providing the Pseudalert ™ reagents and supplies for this study
- Johnson County Department of Public Health for public service bulletin and consultation
- University of Iowa faculty and staff who kindly provided private spa samples for this study